

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 August 2003 (07.08.2003)

PCT

(10) International Publication Number
WO 03/064388 A2

(51) International Patent Classification: C07D 209/42

(21) International Application Number: PCT/IB03/00691

(22) International Filing Date: 29 January 2003 (29.01.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
02290206.8 30 January 2002 (30.01.2002) EP(71) Applicant (for all designated States except US): LES
LABORATOIRES SERVIER [FR/FR]; 12, place de la
Défense, F-92415 Courbevoie Cedex (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SIMIG, Gyula
[HU/HU]; Hollosy Simon u. 25, H-1126 Budapest (HU).
MEZEI, Tibor [HU/HU]; Borz u. 4, H-1221 Budapest
(HU). PORCS-MAKKAY, Marta [HU/HU]; Bem Jozsef
u. 21, H-2013 Pomaz (HU). MANDI, Atila [HU/HU];
Riado u. 12, H-1026 Budapest (HU).(74) Common Representative: LES LABORATOIRES
SERVIER; Direction Brevets, 12, place de la Défense,
F-92415 Courbevoie Cedex (FR).(81) Designated States (national): AE, AG, AI, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE,
SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ,
VC, VN, YU, ZA, ZM, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE,
ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, SE, SI,
SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).**Published:**— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guidance
Notes on Codes and Abbreviations" appearing at the beginning
of each regular issue of the PCT Gazette.(54) Title: PROCESS FOR THE PREPARATION OF HIGH PURITY PIRINDOPRIL AND INTERMEDIATES USEFUL IN THE
SYNTHESIS(57) Abstract: The invention relates to 1-[2(S)-1(1(S)-(ethoxycarbonyl)butylamino)propionyl]-3a(S,7a(S)octahydroindol-2(S)-car-
boxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl
carbodiimide, and a process for the preparation thereof. The invention also relates to new intermediates of the general Formula III
(wherein R stands for lower alkyl or aryl lower alkyl). The compound of the Formula I - pirindopril - is a known ACE inhibitor.

- 1 -

**PROCESS FOR THE PREPARATION OF HIGH PURITY PERINDOPRIL
AND INTERMEDIATES USEFUL IN THE SYNTHESIS**

This invention relates to a process for the preparation of high purity perindopril and intermediates useful in the synthesis of perindopril.

- 5 More particularly the invention is concerned with a process for the synthesis of high purity perindopril free of certain contaminations, intermediates useful in the synthesis and a process for the preparation of said intermediates.

Perindopril – particularly the *t*-butylamine salt thereof – possesses useful pharmacological properties. The main activity of perindopril is the inhibition of the conversion of the
10 enzyme Angiotensin I (or kininase II) into the octapeptide Angiotensin II; thus it is an ACE inhibitor. The above beneficial effect of perindopril enables the use of this active ingredient in the treatment of cardiovascular diseases, particularly arterial hypertension and cardiac insufficiency.

Perindopril is mentioned the first time in EP 0,049,658. However the synthesis of
15 perindopril is not exemplified.

An industrial scale perindopril synthesis is described in EP 0,308,341. The structures of the compounds of formula I to XII, I', VII' and VIII' are described in the annex. According to this process the compound of the Formula V is reacted with the compound of the Formula II in the presence of dicyclohexyl carbodiimide and 1-hydroxy-benzotriazole, whereafter
20 the benzyl ester of the Formula VI is debenzylated to give perindopril of the Formula I, which is then converted into the salt of the Formula I' by reacting with *t*-butylamine.

The drawback of this process is that the purity of the perindopril thus obtained is not satisfactory and for this reason a series of purification steps is required to provide a product which meets the severe quality requirements of pharmaceutical active ingredients. The
25 reason of said disadvantage is that the coupling reaction of the compounds of the Formulae V and II is carried out in the presence of dicyclohexyl carbodiimide which results in the

- 2 -

formation of a considerable amount of contaminations of the benzyl esters of the Formulae VII and VIII which are transformed by debenzylation into the compounds of the Formulae VII' and VIII'. The removal of said contaminations is encountered with significant difficulties.

- 5 According to unpublished French patent application 01.09839 dihydroindole-2-carboxylic acid or its ester of the general Formula IX (wherein R¹ stands for hydrogen or lower alkyl containing 1-6 carbon atoms) is reacted with a compound of the general Formula X (wherein R² is an amino protecting group) in an organic solvent, in the absence or in the presence of not more than 0.6 mole of 1-hydroxy-benzotriazole, related to 1 mole of the compound of the general Formula IX, and 1-1.2 mole of dicyclohexyl carbodiimide, related to 1 mole of the compound of the Formula IX, subjecting the compound of the general Formula XI thus obtained (wherein R¹ and R² are as stated above) to catalytic hydrogenation and converting the compound of the Formula XII thus obtained (wherein R¹ and R² are as stated above) into perindopril in a known manner.
- 10
- 15 It is the object of the present invention to provide a process for the preparation of high purity perindopril free of contaminations derivable from dicyclohexyl carbodiimide, particularly compounds of the Formulae VII' and VIII'.

The above object is solved with the aid of the process and new intermediates of the present invention.

- 20 According to an aspect of the present invention there is provided 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide.

- 25 According to a particular embodiment of the above aspect of the present invention there is provided 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of compounds of the Formula VII' and VIII'.

- 3 -

According to a further aspect of the present invention there is provided a process for the preparation of the compound of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide, particularly free of compounds of the Formula VII' and VIII' which comprises reacting the compound of the Formula II with a suitable carbonic acid derivative; activating the compound of the general Formula III thus obtained (wherein R stands for lower alkyl or aryl-lower alkyl) with thionyl chloride; reacting the activated compound thus obtained with a compound of the Formula IV and if desired reacting the compound of the Formula I thus obtained with *t*-butylamine.

According to a still further aspect of the present invention there are provided compounds of the general Formula III (wherein R stands for lower alkyl or aryl-lower alkyl).

According to a still further aspect of the present invention there is provided a process for the preparation of compounds of the general Formula III (wherein R stands for lower alkyl or aryl-lower alkyl) which comprises reacting the compound of the Formula II with a suitable carbonic acid derivative.

According to a still further aspect of the present invention there are provided pharmaceutical compositions comprising 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide as active ingredient in admixture with suitable inert pharmaceutical carriers.

According to a still further aspect of the present invention there is provided the use of 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide as pharmaceutical active ingredient, particularly as ACE inhibitor.

- 4 -

According to a still further aspect of the present invention there is provided a method of antihypertensive treatment which comprises administering to the patient in need of such treatment a pharmaceutically active amount of 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide.

Synthesis of a peptide bond normally involves the reaction of a carboxy activated N-protected amino-acid with a carboxy protected amino acid. N-Acyl-, or more specifically N-alkoxycarbonyl amino-acid chlorides (derived from acids of Formula III) represent one of the classical groups of carboxy activated N-protected amino-acid derivatives.

Two methods are known to form a peptide bond starting from N-alkoxycarbonyl amino-acid chlorides (see: Houben-Weyl: Methoden der Organischen Chemie Band XV/II, pp. 355-363):

- reaction with amino-acid esters in organic solvent in the presence of equivalent base,
- reaction with amino acids in alkaline aqueous solution (under Schotten-Baumann conditions).

In the modern practise, amino-acid chlorides are considered as over-reactive species leading to undesired side reactions, therefore alternative carboxy activation methods, e.g. the use of DCC, are preferred (see: R. C. Sheppard: Peptide Synthesis in Comprehensive Organic Chemistry, Vol. 5, pp. 339-352, edited by E. Haslam, Pergamon Press, Oxford, 1994).

The present invention is based on the recognition that the new N-alkoxy(aralkoxy)carbonyl amino-acids of general Formula III can be successfully activated with thionyl chloride and the activated carboxylic acid derivative thus obtained can be preferably used for the acylation of the amino-acid of Formula IV in an organic solvent to form the required compound of Formula I in a single reaction step. This recognition is surprising in several respect: the reaction is carried out in organic solvent in the absence of base and not in alkaline water as suggested by prior art; the "over-reactive" character of the amino acid-

- 5 -

chlorides mentioned above does not cause inconvenient side reactions, and the alkoxy(aralkoxy)carbonyl protecting group is removed during the peptide formation process. This recognition is so much the more surprising as in all the actually disclosed syntheses of perindopril the carbonic acid derivative was always activated with dicyclohexyl carbodiimide causing the formation of undesired contaminations. The use of thionyl chloride as activating agent is highly advantageous. On the one hand no difficultly removable by-products are formed, while on the other gaseous hydrogen chloride and sulphur dioxide formed in the reaction can be easily eliminated from the reaction mixture.

The definitions used in the patent specification and the claims are to be interpreted as follows.

The term "lower alkyl" relates to straight or branched chain alkyl groups containing 1-6 carbon atoms (e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, secondary butyl, tertiary-butyl, n-pentyl, n-hexyl etc.), preferably methyl, ethyl or tertiary butyl.

The term "aralkyl" relates to alkyl groups as defined above substituted by one or two aryl groups (e.g. benzyl, beta-phenyl-ethyl, beta-beta-diphenyl-ethyl etc.; preferably benzyl).

According to the first step of the perindopril synthesis according to the present invention the compound of the Formula II is acylated with a suitable carbonic acid derivative. The process is carried out in a manner known per se. As carbonic acid derivative preferably the corresponding alkyl chloroformates or di-alkyl-dicarbonates can be used. The methoxycarbonyl, ethoxycarbonyl and benzyloxycarbonyl group may be preferably introduced with the aid of methyl chloroformate, ethyl chloroformate and benzyl chloroformate, respectively. The tertiary butoxycarbonyl group may be introduced preferably by using di-tertiary butyl dicarbonate. The acylation reaction is carried out in an inert organic solvent and in the presence of a base. As inert organic solvent preferably halogenated aliphatic hydrocarbons (e.g. dichloromethane, dichloroethane or chloroform), esters (e.g. ethyl acetate) or ketones (e.g. acetone), can be used. One may preferably use acetone as solvent. The reaction may be carried out in the presence of an inorganic or organic base. As inorganic base preferably alkali carbonates (e.g. sodium carbonate or

- 6 -

potassium carbonate) or alkali hydrogen carbonates (e.g. sodium hydrogen carbonate or potassium hydrogen carbonate) can be used. As organic base preferably trialkyl amines (e.g. triethylamine) or pyridine can be used. One may preferably use an alkali carbonate or triethylamine as base.

- 5 The reaction is preferably carried out at temperature between 0°C and 30°C, particularly at ambient temperature. One may proceed preferably by preparing the reaction mixture at a low temperature of about 0-5°C, then allowing the reaction mixture to warm to ambient temperature and carrying out the reaction at this temperature for a period of some hours.

- 10 The reaction mixture is worked up in the usual manner. One may proceed preferably by treating the reaction mixture after evaporation with an acid, extracting the mixture with an organic solvent, extracting the organic phase with an aqueous alkali hydroxide solution, acidifying the aqueous layer and extracting the compound of the general Formula III obtained into an organic solvent. The crude product obtained on evaporating the organic phase can be directly used for the further reaction without any purification.

- 15 The compounds of the general Formula III thus obtained are new and are also subject matter of the present invention.

- 20 According to the next step of the perindopril synthesis of the present invention the compound of the general Formula III is activated with thionyl chloride. In this step thionyl chloride is used in an excess, preferably in a molar ratio of 1.1-2 – particularly 1.5-1.7 – related to 1 mole of the compound of the general Formula III. The reaction is carried out in an inert organic solvent. As reaction medium preferably halogenated aliphatic hydrocarbons (e.g. dichloromethane, dichloroethane or chloroform), esters (e.g. ethyl acetate) or ethers (e.g. diethyl ether, tetrahydrofuran, dioxane) can be used. One may carry out the reaction particularly advantageously in dichloro- methane as medium. The
- 25 reaction is carried out at the temperature between 0°C and 30°C, particularly at ambient temperature. One may proceed preferably by preparing the reaction mixture at a lower temperature of 0-5°C, thereafter allowing the reaction mixture to warm to ambient temperature and carrying out the reaction at this temperature for a few hours. The

- 7 -

activation with thionyl chloride being completed the excess of thionyl chloride is removed together with the hydrochloric acid and sulphur dioxide.

5 The activated compound obtained from the compound of the general Formula III is reacted with a compound of the Formula IV. The compound of the Formula IV is perhydroindole-2-carboxylic acid. The reaction is carried out in an inert organic solvent. As a reaction medium preferably a halogenated aliphatic hydrocarbon (e.g. dichloromethane, dichloroethane or chloroform), ester (e.g. ethyl acetate) or ether (e.g. ethyl ether, tetrahydrofuran or dioxane) can be used. One may carry out the reaction preferably in tetrahydrofuran or dichloromethane as medium. The reaction is carried out under heating, 10 preferably at the boiling point of the reaction mixture, advantageously under reflux. The reaction takes place within a few hours. The compound of the Formula IV is used in an amount of 0.5-0.9, preferably 0.7-0.8 moles, related to 1 mole of compound of the general Formula III. The acylation reaction being completed the reaction mixture is concentrated in vacuo.

15 Perindopril of the Formula I may be converted into the *t*-butylamine salt of the Formula I' by reaction with *t*-butylamine. Salt formation may be carried out in a manner known per se. The salt formation reaction is carried out in an inert organic solvent, preferably ethyl acetate. *t*-Butylamine is used preferably in approximately equimolar amount.

20 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide can be used in therapy in the form of pharmaceutical compositions. The preparation of said pharmaceutical compositions, the dosage forms and the daily dosage scheme are similar to those described in prior art for the formulation and pharmaceutical use of perindopril.

25 The starting material of the Formula II is described in EP 0,308,340, EP 0,308,341 and EP 0,309,324. The acid of the Formula IV is described in EP 0,308,339 and EP 0,308,341.

The advantage of the present invention is that it provides highly pure perindopril free of

- 8 -

contaminations derivable from dicyclohexyl carbodiimide. The process is simple and can be easily scaled up. The particular advantage of the present invention is that the use of dicyclohexyl carbodiimide is completely eliminated and therefore there is not even a theoretical possibility of the formation of contaminations which may be derived from dicyclohexyl carbodiimide. Further advantage: the acylation can be carried out with the acid of the Formula IV. Protection of the carboxylic group is not required.

Further details of the present invention are to be found in the following Examples without limiting the scope of protection to said Examples.

Examples

10 **Preparation of *N*-[2-(ethoxycarbonyl)butyl]-*N*-alkoxycarbonylalanine**

Example 1

***N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine**

To a suspension of *N*-[2-(ethoxycarbonyl)butyl]alanine (21.7 g, 100 mmol) in acetone (250 mL) was added a solution of triethylamine (27.7 mL, 20.2 g, 200 mmol) in acetone (50 mL) followed by ethyl chloroformate (24.8 mL, 28.2 g, 260 mmol) at 0-5 °C. After stirring for 2 h at ambient temperature the solvent was evaporated and the residue was stirred with a mixture of water (200 mL) and concentrated hydrochloric acid (2 mL) for 8 h at ambient temperature. The mixture was extracted with ethyl acetate (200 mL) and the solution in ethyl acetate was extracted with cold aqueous sodium hydroxide solution [prepared from ice (100 g) and aqueous sodium hydroxide solution (1N, 200 mL)]. Concentrated hydrochloric acid (15 mL) was added to the aqueous layer and the mixture was extracted with ethyl acetate (200 mL). After drying and evaporation *N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine (23.8 g, 82 %) was obtained as a yellow oil which can be used in the next reaction without further purification.

25 **IR** (film): 3500-2400 (OH st), 1709 (C=O st), ~1200 (C-O st), 898 (OH out of plane b), cm⁻¹.

¹H-NMR (CDCl₃, TMS, 400 MHz): δ 10.14 (1H, bs, COOH), 4.99 and 4.58 (1H, dd, J=4.5, 10.0 Hz, N-CH-CH₂-CH₂-CH₃), 4.29 (2H, q, J=7.1 Hz, N-C(O)-O-CH₂-CH₃), 4.24-

- 9 -

4.13 (2H, m, CH-C(O)-O-CH₂-CH₃), 3.84 (1H, q, J=6.9 Hz, CH₃-CH), [2.08-1.97 (1H, m) and 1.74 -1.62 (1H, m) N-CH-CH₂-CH₂-CH₃], 1.56-1.42 (2H, m, N-CH-CH₂-CH₂-CH₃), 1.51 (3H, d, J=6.9 Hz, CH₃-CH), 1.33 (3H, t, J=7.2 Hz, CH-C(O)-O-CH₂-CH₃), 1.27 (3H, t, J=7.1 Hz, N-C(O)-O-CH₂-CH₃), 0.99 (3H, t, J=7.3 Hz, N-CH-CH₂-CH₂-CH₃).

5 ¹³C-NMR (CDCl₃, TMS, 400 MHz): δ 176.4 and 175.8, 172.5, 155.8, 63.0, 62.7, 58.9, 53.8, 31.4, 19.8, 16.6, 14.0, 13.9, 13.5. (signals of the main rotamer)

Example 2

***N*-[2-(ethoxycarbonyl)butyl]-*N*-methoxycarbonylalanine**

To a suspension of *N*-[2-(ethoxycarbonyl)butyl]alanine (4.35 g, 20 mmol) in acetone
10 (50 mL) was added a solution of triethylamine (5.5 mL, 4.05 g, 40 mmol) in acetone (10 mL) followed by methyl chloroformate (4.0 mL, 4.91 g, 52 mmol) at 0-5 °C. After stirring for 2 h at ambient temperature the solvent was evaporated and the residue was stirred with a mixture of water (40 mL) and concentrated hydrochloric acid (0.4 mL) for 8 h at ambient temperature. The mixture was extracted with ethyl acetate (40 mL) and the
15 solution in ethyl acetate was extracted with cold aqueous sodium hydroxide solution [prepared from ice (20 g) and aqueous sodium hydroxide solution (1N, 40 mL)]. Concentrated hydrochloric acid (3 mL) was added to the aqueous layer and the mixture was extracted with ethyl acetate (40 mL). After drying and evaporation *N*-[2-(ethoxycarbonyl)butyl]-*N*-methoxycarbonylalanine (3.84 g, 70 %) was obtained as a
20 yellow oil which can be used in the next reaction without further purification.

IR (film): ~ 3400 (OH st), 1713 (C=O st), 1294 (OH in plane b), 1205 (C-O st ester) cm⁻¹.

¹H-NMR (CDCl₃, TMS, 400 MHz): δ ~10.2 (1H, bs, COOH), 5.00 and 4.60 (1H, s, CH-CH₂-CH₂-CH₃), 4.28 (2H, q, J=7.2 Hz, O-CH₂-CH₃), 3.96 and 3.85 (1H, m, CH-CH₃), 3.75 (3H, s, OCH₃), [2.10-1.98 (1H, m) and 1.74-1.62 (1H, m) CH-CH₂-CH₂-CH₃], 1.56-1.42 (2H, m, CH-CH₂-CH₂-CH₃), 1.50 (3H, d, J=6.9 Hz, CH-CH₃), 1.33 (3H, t, J=7.2 Hz, O-CH₂-CH₃), 0.99 (3H, t, J=7.3 Hz, CH-CH₂-CH₂-CH₃).

25

Example 3

***N*-[2-(ethoxycarbonyl)butyl]-*N*-*i*-butyloxycarbonylalanine**

- 10 -

To a suspension of *N*-[2-(ethoxycarbonyl)butyl]alanine (1.1 g, 5 mmol) and potassium carbonate (0.76 g, 5.5 mmol) in acetone (15 mL) was added di-*t*-butyl dicarbonate (1.20 g, 5.5 mmol) and water (1.25 mL) at 0-5 °C. After stirring for 2 h at ambient temperature the mixture was cooled and the solid precipitation was filtered off. Ethyl acetate (15 mL), ice (5 g) and aqueous NaOH solution (1N, 10 mL) was added. After stirring for 5 min the layers were separated. Concentrated hydrochloric acid (1.5 mL) was added to the aqueous layer and the mixture was extracted with ethyl acetate (15 mL). After drying and evaporation *N*-[2-(ethoxycarbonyl)butyl]-*N*-*t*-butyloxycarbonylalanine (0.61 g, 38 %) was obtained as a yellow oil which can be used in the next reaction without further purification.

IR (KBr): ~ 3400 (OH st), 1705 (C=O st acid), 1299 (OH in plane b) ~ 1690 (C=O st amide), 1771 (C=O st ester), 1159 (C-O st ester) cm⁻¹.

¹H-NMR (CDCl₃, TMS, 400 MHz): δ ~9.2 (1H, bs, COOH), 5.00 (1H, dd, J=4.4, 10.2 Hz, CH-CH₂-CH₂-CH₃), 4.29 (2H, q, J=7.1 Hz, O-CH₂-CH₃), 3.69 (1H, q, J=6.8 Hz, CH-CH₃), [2.10-1.98 (1H, m) and 1.70-1.56 (1H, m), CH-CH₂-CH₂-CH₃], 1.56-1.42 (2H, m, CH-CH₂-CH₂-CH₃), 1.49 (3H, d, J=6.8 Hz, CH-CH₃), 1.46 (9H, s, C(CH₃)₃), 1.34 (3H, t, J=7.1 Hz, O-CH₂-CH₃), 0.99 (3H, t, J=7.3 Hz, CH-CH₂-CH₂-CH₃).

Example 4

***N*-[2-(ethoxycarbonyl)butyl]-*N*-benzyloxycarbonylalanine**

To a suspension of *N*-[2-(ethoxycarbonyl)butyl]alanine (2.2 g, 10 mmol) and potassium carbonate (2.2 g, 16 mmol) in a mixture of acetone (30 mL) and water (2.5 mL) was added benzyl chloroformate (2.0 mL, 2.4 g, 14 mmol) at 0-5 °C. After stirring for 2 h at ambient temperature the solid was filtered off, the solvent was evaporated, the residue was stirred with cold aqueous sodium hydroxide solution [prepared from ice (20 g) and aqueous sodium hydroxide solution (1N, 40 mL)] and extracted with ethyl acetate (40 mL). The aqueous layer was acidified with an aqueous solution of hydrochloric acid 1/1 (20 mL) and the mixture was extracted with ethyl acetate (40 mL). After drying and evaporation *N*-[2-(ethoxycarbonyl)butyl]-*N*-benzyloxycarbonylalanine (1.66 g, 47%) was obtained as a yellow oil which can be used in the next reaction without further purification.

IR (film): ~ 3400 (OH st), 1710 (C=O st), 699 (CH arom) cm⁻¹.

- 11 -

¹H-NMR (CDCl₃, TMS, 400 MHz): δ ~ 9.0 (1H, bs, COOH), 5.00 (5H, m, Ph), 5.2-5.1 (2H, m, Ph-CH₂), 4.96 and 4.57 (1H, dd, J=9.6, 4.8, N-CH-CH₂-CH₂-CH₃), 4.26 and 4.12 (2H, q, J=7.1 Hz, O-CH₂-CH₃), 4.02 and 3.89 (1H, q, J=6.9 Hz, CH-CH₃), [2.10-1.93 (1H, m) and 1.72-1.62 (1H, m), CH-CH₂-CH₂-CH₃], 1.54 and 1.48 (3H, d, J=7.0 Hz, CH-CH₃), 1.56-1.42 (2H, m, CH-CH₂-CH₂-CH₃), 1.31 and 1.22 (3H, t, J=7.2 Hz, O-CH₂-CH₃), 0.98 and 0.93 (3H, t, J=7.3 Hz, CH-CH₂-CH₂-CH₃).

Preparation of perindopril eburmine

Example 5

Acylation of perhydroindole-2-carboxylic acid using *N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine

To a solution of *N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine (10.1 g, 35 mmol) in dichloromethane (35 mL) thionyl chloride (4.2 mL, 6.9 g, 58 mmol) was added in drops at 0-5°C. It was stirred at ambient temperature for 2-3 h. The solvent was evaporated to give a reddish oil. The residue was dissolved in THF (37.5 mL) and it was added to a suspension of perhydroindole-2-carboxylic acid (4.7 g, 28 mmol) in THF (37.5 mL). The suspension was refluxed with stirring for 4-4.5 h until a brownish solution was formed. After evaporation of the solvent the residue was dissolved in ethyl acetate (120 mL), *t*-butylamine (2.8 mL, 1.95 g, 27 mmol) in ethyl acetate (60 mL) was added slowly to the stirred solution resulting in separation of a crystalline mass. The mixture was heated until a solution was formed, then treated with charcoal. The crystalline product obtained after cooling was filtered to give perindopril eburmine (6.8 g, 55 %).

Example 6

Acylation of perhydroindole-2-carboxylic acid using *N*-[2-(ethoxycarbonyl)butyl]-*N*-methoxycarbonylalanine

Perindopril eburmine was prepared analogously to *Example 5*, using *N*-[2-(ethoxycarbonyl)butyl]-*N*-methoxycarbonylalanine (3.4 g, 12.5 mmol) and perhydroindole-2-carboxylic acid (1.7 g, 10 mmol). The crystalline product obtained was filtered to give perindopril eburmine (2.4 g, 54 %).

Example 7**Acylation of perhydroindole-2-carboxylic acid using *N*-[2-(ethoxycarbonyl)butyl]-*N*-*t*-butoxycarbonylalanine**

Perindopril eburmine was prepared analogously to *Example 5*, using *N*-[2-(ethoxycarbonyl)butyl]-*N*-*t*-butoxycarbonylalanine (0.69 g, 2.2 mmol) and perhydroindole-2-carboxylic acid (0.29 g, 1.7 mmol). The crystalline product obtained was filtered to give perindopril eburmine (0.37 g, 49 %).

Example 8**Acylation of perhydroindole-2-carboxylic acid using *N*-[2-(ethoxycarbonyl)butyl]-*N*-benzyloxycarbonylalanine**

Perindopril eburmine was prepared analogously to *Example 5*, using *N*-[2-(ethoxycarbonyl)butyl]-*N*-benzyloxycarbonylalanine (1.41 g, 4 mmol) and perhydroindole-2-carboxylic acid (0.51 g, 3 mmol). The crystalline product obtained was filtered to give perindopril eburmine (0.60 g, 45 %).

Example 9**Acylation of perhydroindole-2-carboxylic acid using *N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine**

To a solution of *N*-[2-(ethoxycarbonyl)butyl]-*N*-ethoxycarbonylalanine (1.45 g, 5 mmol) in dichloromethane (7.5 mL) thionyl chloride (0.6 mL, 0.98 g, 8.5 mmol) was added dropwise at 0-5°C. It was stirred at ambient temperature for 2-3 h. The excess of thionyl chloride and the sulphur dioxide and hydrogen chloride formed was eliminated in slight vacuo. To the dichloromethane solution thus obtained was added perhydroindole-2-carboxylic acid (0.71 g, 4.2 mmol) and dichloromethane (5.0 mL). The suspension was refluxed with stirring for 2 h until a brownish solution was formed. After evaporation of the solvent the residue was dissolved in ethyl acetate (20 mL), whereupon *t*-butylamine (0.42 mL, 0.29 g, 4.05 mmol) in ethyl acetate (5.0 mL) was added slowly to the stirred

- 13 -

solution resulting in separation of a crystalline mass. The mixture was heated until a solution was formed, then treated with charcoal. The crystalline product obtained after cooling was filtered to give perindopril ebumine (0.64 g, 35 %).

What we claim is,








1. 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide.
- 5 2. 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' according to Claim 1 free of compounds of the Formula VII' and VIII'.
3. Process according to Claim 1 or 2 for the preparation of the compound of the Formula I and the *t*-butylamine salt of the Formula I' free of contaminations derivable
10 from dicyclohexyl carbodiimide, particularly free of compounds of the Formula VII' and VIII' which comprises reacting the compound of the Formula II with a suitable carbonic acid derivative; activating the compound of the general Formula III thus obtained (wherein R stands for lower alkyl or aryl-lower alkyl) with thionyl chloride; reacting the activated compound thus obtained with the compound of the Formula IV and if desired reacting the
15 compound of the Formula I thus obtained with *t*-butylamine.
4. Process according to Claim 3 which comprises using as carbonic acid derivative methyl chloroformate, ethyl chloroformate, benzyl chloroformate or di-*t*-butyl dicarbonate.
5. Process according to any of Claims 3 to 4 which comprises carrying out acylation of the compound of the Formula II in the presence of a base, preferably an alkali
20 carbonate, alkali hydrogen carbonate or tertiary amine.
6. Process according to Claim 5 which comprises using sodium carbonate, potassium carbonate or triethylamine as base.
7. Process according to any of Claims 3 to 6 which comprises carrying out the reaction in an inert organic solvent.

8. Process according to Claim 7 which comprises using a halogenated aliphatic hydrocarbon, aromatic hydrocarbon, ester or ketone as solvent.
9. Process according to Claim 8 which comprises using acetone as solvent.
- 5 10. Process according to Claim 3 which comprises using thionyl chloride in a molar ratio of 1.1-2 preferably 1.5-1.7, related to 1 mole of the compound of the general Formula III.
11. Process according to Claim 10 which comprises carrying out activation of the compound of the general Formula III with thionyl chloride in an inert organic solvent.
- 10 12. Process according to Claim 11 which comprises using a halogenated aliphatic hydrocarbon, ester or ether as inert organic solvent.
13. Process according to Claim 12 which comprises using dichloromethane.
14. Process according to Claim 3 which comprises carrying out the reaction with the compound of the Formula IV in an inert organic solvent.
- 15 15. Process according to Claim 14 which comprises using a halogenated aliphatic hydrocarbon, ester or ether as inert organic solvent.
16. Process according to Claim 15 which comprises using tetrahydrofuran or dichloromethane.
17. Compounds of the general Formula III (wherein R stands for lower alkyl or aryl-lower alkyl).
- 20 18. Compounds according to Claim 17 wherein R stands for methyl, ethyl, tertiary butyl or benzyl.

19. Process for the preparation of compounds of the general Formula III (wherein R stands for lower alkyl or aryl-lower alkyl) which comprises reacting the compound of the Formula II with a suitable carbonic acid derivative.
20. Process according to Claim 19 which comprises using as carbonic acid derivative methyl chloroformate, ethyl chloroformate, benzyl chloroformate or di-*t*-butyl dicarbonate.
21. Process according to any of Claims 19 or 20 which comprises carrying out acylation of the compound of the Formula II in the presence of a base, preferably an alkali carbonate, alkali hydrogen carbonate or tertiary amine.
22. Process according to Claim 21 which comprises using sodium carbonate, potassium carbonate or triethylamine as base.
23. Process according to any of Claims 19 to 22 which comprises carrying out the reaction in an inert organic solvent.
24. Pharmaceutical compositions comprising 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide as active ingredient in admixture with suitable inert pharmaceutical carriers.
25. Pharmaceutical compositions according to Claim 24 as ACE inhibitors.
26. Use of 1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I and the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide as pharmaceutical active ingredient, particularly as ACE inhibitor.

- 17 -

27. Method of antihypertensive treatment which comprises administering to the patient in need of such treatment a pharmaceutically active amount of (1-{2(S)-[1(S)-(ethoxycarbonyl)butylamino]propionyl}-(3aS,7aS)octahydroindol-2(S)-carboxylic acid of the Formula I or the *t*-butylamine salt of the Formula I' thereof free of contaminations derivable from dicyclohexyl carbodiimide.
- 5



2/2

